

Dissecting the instant blood-mediated inflammatory reaction in islet xenotransplantation

Goto M, Tjernberg J, Dufrane D, Elgue G, Brandhorst D, Ekdahl KN, Brandhorst H, Wennberg L, Kurokawa Y, Satomi S, Lambris JD, Gianello P, Korsgren O, Nilsson B. Dissecting the instant blood-mediated inflammatory reaction in islet xenotransplantation. *Xenotransplantation* 2008; 15: 225–234. © 2008 Wiley Periodicals, Inc.

Abstract: Background: A massive destruction of transplanted tissue occurs immediately following transplantation of pancreatic islets from pig to non-human primates. The detrimental instant blood-mediated inflammatory reaction (IBMIR), triggered by the porcine islets, is a likely explanation for this tissue loss. This reaction may also be responsible for mediating an adaptive immune response in the recipient that requires a heavy immunosuppressive regimen.

Materials and methods: Low molecular weight dextran sulfate (LMW-DS) and the complement inhibitor Compstatin were used in a combination of in vitro and in vivo studies designed to dissect the xenogeneic IBMIR in a non-human primate model of pancreatic islet transplantation. Adult porcine islets (10 000 IEQs/kg) were transplanted intraportally into three pairs of cynomolgus monkeys that had been treated with LMW-DS or heparin (control), and the effects on the IBMIR were characterized. Porcine islets were also incubated in human blood plasma in vitro to assess complement inhibition by LMW-DS and Compstatin.

Results: Morphological scoring and immunohistochemical staining revealed that the severe islet destruction and macrophage, neutrophilic granulocyte, and T-cell infiltration observed in the control (heparin-treated) animals were abrogated in the LMW-DS-treated monkeys.

Both coagulation and complement activation were significantly reduced in monkeys treated with LMW-DS, but IgM and complement fragments were still found on the islet surface. This residual complement activation could be inhibited by Compstatin in vitro.

Conclusions: The xenogeneic IBMIR in this non-human primate model is characterized by an immediate binding of antibodies that triggers deleterious complement activation and a subsequent clotting reaction that leads to further complement activation. The effectiveness of LMW-DS (in vivo and in vitro) and Compstatin (in vitro) in inhibiting this IBMIR provides the basis for a protocol that can be used to abrogate the IBMIR in pig-human clinical islet transplantation.

Masafumi Goto,¹ Jenny Tjernberg,² Denis Dufrane,³ Graciela Elgue,² Daniel Brandhorst,² Kristina Nilsson Ekdahl,² Heidi Brandhorst,² Lars Wennberg,⁴ Yoshimochi Kurokawa,⁵ Susumu Satomi,⁶ John D. Lambris,⁷ Pierre Gianello,³ Olle Korsgren² and Bo Nilsson²

¹Tohoku University International Advanced Research and Education Organization, Tohoku University, Sendai, Japan, ²Department of Radiology, Oncology and Clinical Immunology, Division of Clinical Immunology, The Rudbeck Laboratory, University Hospital, Uppsala, Sweden, ³Université Catholique de Louvain, Faculty of Medicine, Experimental Surgery Unit, Brussels, Belgium, ⁴Department of Transplantation Surgery, Karolinska Institute, Huddinge University Hospital, Stockholm, Sweden, ⁵Tohoku University Innovation of New Biomedical Engineering Center, Tohoku University, Sendai, Japan, ⁶Division of Advanced Surgical Science and Technology, Tohoku University, Sendai, Japan, ⁷Department of Pathology and Laboratory Medicine, University of Pennsylvania, Medical School, Philadelphia, PA, USA

Key words: adult porcine islet – compstatin – instant blood-mediated inflammatory reaction – low molecular weight dextran sulfate – xenotransplantation

Address reprints request to Bo Nilsson, Department of Radiology, Oncology and Clinical Immunology, Division of Clinical Immunology, The Rudbeck Laboratory, University Hospital, Uppsala, Sweden (E-mail: bo.nilsson@klinimm.uu.se)

Received 27 April 2008;
Accepted 29 May 2008

Introduction

Clinical islet transplantation is a promising treatment for type I diabetic patients. The improved protocol introduced by Shapiro et al.

in 2000 [1] has greatly improved the results of this approach, but despite these advances, islets derived from more than one donor pancreas are still generally required to cure an individual diabetic patient. This requirement has drawn

attention to the limited availability of human islets for transplantation and sparked interest in the use of islets from alternative sources, particularly the pig [2].

One obstacle to be surmounted before porcine islets can be used in clinical islet xenotransplantation is the injurious instant blood-mediated inflammatory reaction (IBMIR) that elicits massive cell destruction when porcine islets are exposed to fresh human blood [3]. The xenogeneic IBMIR is characterized by activation of platelets and the coagulation and complement systems. This activation is accompanied by infiltration of the islets by polymorphonuclear lymphocytes (PMNs) [3].

The occurrence of this deleterious IBMIR is supported by studies demonstrating that porcine islets are immediately destroyed when transplanted intraportally into the liver of non-human primates [4,5]. Kirchof et al. [6] reported that most of their porcine islet xenografts (22–73%) were substantially damaged after 24 h when transplanted into non-immunosuppressed monkeys. The grafts exhibited cell destruction, with deposition of coagulation and complement components and platelets, supporting the contention that the IBMIR contributes to the islet damage in this model. Further support for the importance of the IBMIR comes from the observation that although porcine islets can successfully survive in the liver of diabetic monkeys for more than 100 days [7,8], very high quantities of islets (25 000 and 50 000 IEQs/kg BW, respectively) are needed to produce normoglycemia in the monkeys, indicating that there is a substantial loss of transplanted tissue.

Using in vitro and small-animal models, we have previously demonstrated that low molecular weight dextran sulfate (LMW-DS) effectively inhibits the activation of the coagulation and complement systems and the infiltration of leukocytes into the islets during xenogeneic islet transplantation [9]. In this study, we have used LMW-DS together with Compstatin, a new peptide complement inhibitor that is suitable for use in clinical islet xenotransplantation [10], to dissect the IBMIR in in vivo (LMW-DS) and in in vitro (LMW-DS and Compstatin) xenotransplantation models. The results of these studies have broadened our understanding of the innate immune events that might be expected to occur in clinical islet xenotransplantation and have provided the basis for a protocol for abrogating the IBMIR during clinical transplantation with porcine pancreatic islets.

Materials and methods

Animals

Retired breeder pigs, weighing approximately 200 kg, were used as donors for all experiments. Cynomolgus monkeys (*Macaca fascicularis*; 3- to 6-yr old; 4–6 kg) were used as recipients. All procedures using pigs were approved by the Swedish Council on Medical Ethics. Cynomolgus monkeys were housed according to the guidelines of the Belgian Ministry of Agriculture and Animal Care. All procedures using monkeys were approved by the local Ethical Committee for Animal Care of the Université Catholique de Louvain.

Islet isolation

Isolation of porcine islets was performed as previously described [11], with minimal modifications. Purified islet fractions were pooled and cultured at 37 °C in a humidified atmosphere with 5% CO₂ in CMRL 1066 medium (Biochrom, Berlin, Germany) supplemented with 20% heat inactivated porcine serum, 2 mM *N*-acetyl-L-alanyl-L-glutamine, 10 mM *N*-2-hydroxyethylpiperazine-N1-2-ethanesulfonic acid (HEPES), 100 IU/ml penicillin, 100 µg/ml streptomycin (Biochrom), and 20 µg/ml ciprofloxacin (Bayer, Leverkusen, Germany).

Evaluation of porcine islet quality

The in vitro function and viability of the porcine islets were assessed after overnight culture as described above. Islet viability determined by trypan blue exclusion assay and insulin release defined as the ratio of stimulated (16.5 mM glucose) to basal (1.65 mM glucose) insulin release, were performed as previously described [11]. For assays of islet insulin content, 1-ml samples were washed with distilled water, then sonicated (Labsonic; Braun, Melsungen, Germany) for 30 s. A 200-µl aliquot of each sample was subjected to acid-ethanol extraction (0.18 M HCl) and used for insulin measurement. Another 100-µl aliquot was dried at 60 °C overnight for consecutive fluorometric DNA assays [12], using calf thymus DNA type I (Sigma, Deisenhofen, Germany) as a standard; 24-h insulin secretion: Immediately after a medium change, 500-µl samples of the medium were taken in duplicate from the remaining Petri dishes for determination of insulin accumulation in the medium, to calculate the 24-h insulin secretion by the islets. Transplantation of islets into nude mice was performed as previously described [11].

Islet transplantation

Before each experiment, the monkeys were sedated with 6 mg/kg Zoletil® 100 (Virbac S.A., Carros, France) intramuscularly, and general anesthesia was maintained with inhalation of 1–3% enflurane. During the experiment, electrocardiogram, blood pressure, and pulse were continuously monitored. The pig islets were suspended in 10 ml of transplant medium (Ringer acetate; Braun) with 25% (w/v) human albumin and 5 mM glucose and injected slowly into the portal vein over the course of 5 min. The animals were treated in pairs, with each pair being given porcine islets from the same donor. One recipient in each pair received LMW-DS (monkeys M5, M7, and M9) and the other heparin as a control (monkeys M6, M8, and M10):

1. Intravenous infusion of LMW-DS (MW 5000; Sigma Chemicals, St Louis, MO, USA) was performed via an indwelling catheter placed in the jugular vein or via a catheter in the portal vein. In the LMW-DS-treated groups, dextran with a molecular weight of 1 kDa (Promiten, Pharmalink AB, Upplands Väsby, Sweden) was injected i.v. just before islet transplantations to avoid the risk of anaphylactoid reactions triggered by LMW-DS. After the injection of Promiten, the monkey received a bolus dose of LMW-DS (1.5 mg/kg) i.v. prior to islet infusion, followed by 3.0 mg/kg LMW-DS given together with the porcine islets (10 000 IEQs/kg of recipient BW). The transplantation was followed by a continuous i.v. infusion of LMW-DS (1.0–1.5 mg/kg/h) for up to 24 h.
2. In the heparin-treated groups, the monkeys received a continuous i.v. infusion of heparin (35 U/kg of BW, heparin LEO, 5000 U/ml; LEO Pharma Nordic, Malmö, Sweden) for 24 h, beginning immediately prior to islet infusion.

Blood samples

All blood samples from the monkeys were drawn from a femoral vein catheter at 0, 15, 30, 45, 60, 75, 90, 120 min and 24 h after transplantation. Blood was also drawn from healthy human blood donors into 7-ml tubes containing citrate, EDTA, or 500 µg of hirudin, a specific inhibitor of thrombin (Refludan; Pharmion Ltd, Cambridge, UK). To obtain plasma, the samples were centrifuged at 4500 g for 5 min. If not immediately analyzed for activated partial thromboplastin time (APTT), the samples were stored at 70 °C.

Analyses of blood and plasma samples

Activated partial thromboplastin time measurements were performed as previously described [13]. Platelet counts and differential leukocyte counts were obtained using a Coulter-AcT-diff analyzer (Beckman Coulter, Miami, FL, USA) and EDTA-treated blood. Plasma levels of thrombin–anti-thrombin (TAT) were quantified using commercially available EIA kits (TAT; Behringwerke, Marburg, Germany). C3a generation was measured in plasma according to the method of Nilsson Ekdahl et al. [14], and sC5b–9 was analyzed using a modification of the enzyme immunoassay described by Nilsson Ekdahl et al. [14] and Mollnes et al. [15].

Plasma interleukin-6 (IL-6), tumor necrosis factor- α (TNF α), IL-1 β , and C-reactive protein (CRP) were measured using a commercial ELISA kit (Immulite IL-6, Immulite TNF α , Immulite IL-1 β , and Immulite High Sensitivity CRP, respectively; Diagnostic Products Corporation, Los Angeles, CA, USA).

Histological and immunohistochemical staining

The monkey livers bearing transplanted adult porcine islet grafts were retrieved 24 h after transplantation, at a time when the major part of the IBMIR has generally occurred [3]. Some tissue samples were snap-frozen in isopentane and stored at –70 °C. Other samples were fixed with 4% *p*-formaldehyde overnight, and then embedded in paraffin. The samples were sectioned and subsequently used for morphological scoring after hematoxylin eosin staining.

Immunohistochemical staining was carried out using guinea pig anti-insulin (DAKO, Carpinteria, CA, USA), mouse anti-human neutrophil elastase (DAKO), mouse anti-human CD68 (DAKO), mouse anti-human MAC 387 (Abcam, Cambridge, UK), mouse anti-human CD56 (Monosan, Stockholm, Sweden), rabbit anti-human CD3 (DAKO), mouse anti-human CD20 (DAKO), rabbit anti-human IgG and IgM (DAKO), mouse anti-human CD41 (DAKO), mouse anti-human C3c (QUIDEL, San Diego, CA, USA), or goat anti-human C9 (Serotec Ltd Scandinavia, Oslo, Norway).

Treatment of porcine islets with human plasma

Approximately 1000 pig islets/40 µl of plasma (typically 5000 islets in 200 µl) were incubated in human hirudin-treated plasma in heparinized test tubes. Five different islet preparations and five

different plasma preparations were used in these experiments. In some experiments, hirudin-treated plasma was pre-incubated with 20 μM (final concentration) of the potent Compstatin analog, Ac-ICV(1-MeW)QDWGAHRCT-NH₂ [16], for 15 min at 37 °C before the islets were added. The mixture of islets and plasma was then incubated, with gentle shaking, at 37 °C for up to 30 min. After centrifugation, the islets were immediately prepared for complex object parametric analyzer and sorter (COPAS) analysis and confocal microscopy.

Preparation of islets for flow cytometry and confocal microscopy

Ten microliters of fluorescein isothiocyanate (FITC)-labeled antibody recognizing one of the following proteins was added to 5000 islets (corresponding to approximately 10×10^6 cells) in 100 μl of phosphate-buffered saline (PBS) according to the manufacturer's recommendations for single cells: C1q (1.0 g/l; AbCam), C3c (3.2 g/l, for detection of C3b and iC3b; DakoCytomation, Glostrup, Denmark), C4 (1.3 g/l; DakoCytomation), C9 (2.6 g/l; DakoCytomation), mannose-binding lectin (MBL) (0.7 g/l; DakoCytomation), IgG (2.6 g/l; DakoCytomation), or IgM (4.0 g/l; DakoCytomation). Irrelevant mouse IgG1 (0.1 g/l; DakoCytomation) was used as a negative control. For all immunostaining experiments, the islets were incubated, while gently rotating on ice, for 30 min in the presence of an individual antibody. After being washed with PBS, the islets were treated with 1% formaldehyde (Apoteket, Gothenburg, Sweden) and kept on ice until analyzed.

Complex object parametric analyzer and sorter analysis

The fluorescence-stained islets were analyzed using a COPAS (Union Biometrica, Somerville, MA, USA), which is a large particle-based flow cytometry instrument [17]. For each experiment, 1000 islets were analyzed using a 488/514 multi-line laser, and positive cells were sorted out for further analysis by confocal microscopy. The COPAS flow cytometry data were analyzed using CellQuest Pro software (BD Biosciences Immunocytometry Systems, San Jose, CA, USA). Data were reported as mean fluorescent intensity (MFI).

Confocal microscopy

One to two hundred hand-picked, stained islets were contained in a drop of PBS in a small Petri dish and protected from light before examination in the confocal microscope (Zeiss 510 Meta con-

focal; Carl Zeiss, Jena, Germany). Examination of the stained islets was performed using the 488-nm laser at 10 times magnification. Counter staining with 4',6-diamidino-2-phenylindole was used to visualize the nuclei of living islet cells.

Complement inhibition assay

One hundred microliters of 10% human serum (v/v), diluted in veronal buffer with 1 mM Ca²⁺, 0.3 mM Mg²⁺, 1% (w/v) bovine serum albumin, and 0.05% (v/v) Tween 20, was incubated in the presence of serially diluted LMW-DS and/or Compstatin in wells of microtiter plates for 30 min at 37 °C. The wells were then washed with PBS containing 0.05% (v/v) Tween 20, and the bound C3 fragments were detected using 100 μl of horseradish peroxidase-conjugated anti-C3c (Dako AS, Glostrup, Denmark).

Statistical analysis

All values are expressed as mean \pm SEM and were compared using Student's unpaired *t*-test or using the Mann-Whitney test for unpaired samples. Values of *P* < 0.05 were considered statistically significant.

Results

Islet quality

The viability of the adult porcine islets (APIs) used in this study was 96, 100, and 97%, respectively. The stimulation index in the static glucose stimulation (SGS) test was 1.29, 1.84, and 1.40, and the mean insulin content was 613, 149, and 685 $\mu\text{U}/\text{IEQs}$, respectively. Adult porcine islets used in each experiment cured diabetic athymic mice. When we assessed the possible detrimental effect of LMW-DS by incubating APIs from three different pancreata in the presence (100, 1000, or 2500 mg/l) or absence of LMW-DS, we found no adverse effect of LMW-DS on insulin release at any of the concentrations tested (data not shown).

Influence of LMW-DS on blood cell counts, liver and renal function, and cytokine induction in transplanted monkeys

One of the transplanted control monkeys (M6) treated with heparin died 2 h after transplantation due to severe hypoglycemia. The platelet and leukocyte counts and the creatinine levels were kept within normal ranges throughout the experiments with one exception: The granulocyte count

tended to increase 2 h after transplantation in the heparin-treated group (3.9 ± 0.5 vs. 9.6 ± 1.6) compared to that of the LMW-DS-treated group (6.0 ± 0.9 vs. 7.3 ± 1.4). There was also a tendency towards an increase in the liver enzymes at 24 h after islet transplantation in the heparin-treated monkeys [heparin vs. LMW-DS: aspartate aminotransferase (AST), 434.7 ± 126.4 vs. 288.0 ± 130.4 ; alanine aminotransferase (ALT), 207.7 ± 68.7 vs. 116.8 ± 47.7]. No bleedings or other adverse reactions were observed.

Influence of LMW-DS on cytokine induction was examined using three healthy monkeys. Only a slight increase in the IL-6 levels was seen 24 h after administration of LMW-DS in two out of three healthy monkeys (maximum $27 \mu\text{g/l}$). However,

LMW-DS did not trigger an increase of plasma IL-1 β , TNF α , or CRP (not shown).

LMW-DS concentrations in transplanted monkeys

Previous studies showed a strong correlation between APTT and the concentration of LMW-DS [13]. Plasma APTT was therefore used to follow the blood concentration of LMW-DS in the transplanted monkeys (Fig. 1). The APTT in monkeys treated with heparin at concentrations routinely used in clinical islet transplantation (i.e. 500–1000 IU/l) was kept constant at 25–40 s throughout the whole study period. The APTT in monkeys treated with LMW-DS reached around 100 s at 15 min after islet infusion, but gradually decreased during 2 h after islet transplantation. After 24 h, the APTTs in monkeys M5, M7, and M9 were 101, 66, and 107 s, respectively. Thus, both M5 and M9 had higher concentrations of LMW-DS compared with M7.

Inhibition of the IBMIR by LMW-DS during pig islet xenotransplantation

Low molecular weight dextran sulfate, unlike heparin, diminished both the coagulation and the complement cascade activation in two sets of monkeys. The increase of coagulation marker TAT was effectively inhibited by LMW-DS (Fig. 2). The complement activation parameters C3a and sC5b-9 were also suppressed by LMW-DS in both treated monkeys compared to the controls during the study period (Fig. 2). In M5, TAT was totally suppressed while C3a was more difficult to evaluate without the corresponding control (M6). In this animals, C5b-9 was not assessed due to an insufficient amount of plasma samples.

Histological evaluation of grafted pig islets after intraportal transplantation into monkeys treated with LMW-DS or heparin

Morphological aspects of islet grafts were scored semi-quantitatively according to the representative examples shown in Fig. 3. As summarized in Table 1, histology of the transplanted grafts were well kept in the monkeys treated with LMW-DS in both settings of experiments. However, the beneficial effects of LMW-DS were more pronounced in M5 and M9 compared with M7. Indeed, the completely preserved islets (score 0 in all categories) were encountered in 37.2 and 44% of the LMW-DS treated animals M5 and M9 (LMW-DS treated monkeys), respectively, but in only 22% of the control M10.




Fig. 1. Plasma APTT values in transplanted diabetic monkeys (M5 and M7–M10) treated with heparin (squares) or LMW-DS (circles).




Fig. 2. EDTA blood was drawn from a femoral vein catheter of the transplanted monkeys treated with heparin (squares) or LMW-DS (circles) at varying time points after porcine islet xenotransplantation. TAT, C3a, and sC5b-9 levels were assessed and expressed as percentage of the pre-transplant values.

Immunohistochemical staining of grafted pig islets after intraportal transplantation into monkeys treated with LMW-DS or heparin

The immunohistochemical findings from the grafts were summarized in Fig. 4 and Table 2. As expected, most parameters involved in innate immune responses were active after 24 h post-islet transplantation in the controls M8 and M10. In particular, CD68+ macrophages, and neutrophil elastase positive PMNs were abrogated in the monkeys treated with LMW-DS compared with the controls given heparin. Also, CD41+ platelets tended to be lower in the LMW-DS treated animals. CD56+ natural killer cells were found only occasionally. Unlike the soluble complement markers there was no clear inhibition of complement activation as reflected in deposition of C3

fragments and C9 on the surface of the islets. Furthermore, IgM antibodies were found on islet both in LMW-DS and heparin-treated animals. Most of parameters reflecting specific immune responses were yet silent. However, CD3+ T-cell infiltration was already seen in the islet grafts of the controls M8 and M10. Notably, this infiltration was effectively suppressed by LMW-DS.

Binding of complement components to porcine islets after incubation in human plasma

After incubation in hirudin-treated plasma, the porcine islets were stained with FITC-conjugated antibodies recognizing IgG, IgM, C1q, C3b/iC3b, C4 fragments, C9, and MBL. Large particle flow cytometry and confocal microscopy demonstrated that antibodies against IgG, IgM, C1q, C4, and C3

Instant blood-mediated inflammatory reaction and islet xenotransplantation




Fig. 3. Visual examples of the morphological scoring system used to quantify different aspects of the IBMIR. Hematoxylin eosin-stained porcine islet grafts retrieved 24 h after intraportal xenotransplantation from diabetic monkeys treated with LMW-DS or heparin were used. A summary of all transplanted monkeys is presented in Table 1.

bound strongly to the islets, but the binding of MBL and C9 was less prominent (Fig. 5A and B). C3b/iC3b fragments were detected on the islets after only 5 min, and the binding of C3b/iC3b continued to increase over time. Addition of Compstatin significantly reduced the binding of C3b/iC3b to the islets (Fig. 5C). Confocal microscopy analyses confirmed these results (not shown).

Inhibition of complement activation by LMW-DS and Compstatin

Ten percentage (v/v) human serum was incubated in wells of microtiter plates in the presence of LMW-DS and/or Compstatin for 30 min at 37 °C (Fig. 6). In the presence of Compstatin there was no effect below 0.5 μM of the compound, but at higher concentrations Compstatin gradually inhibited complement activation. At 5 μM total inhibition was achieved. LMW-DS inhibited complement activation only marginally between 10 and 100 mg/l, but the effect was more pronounced at concentrations above this level. There was no indication of interaction between the drugs regarding this effect on complement activation in serum.

Discussion

We have previously shown that LMW-DS efficiently prevents clotting that occurs in both allogeneic and xenogeneic IBMIR triggered by APIs both in vitro and in vivo in a small animal model [9,18]. Here, we confirm that LMW-DS is efficient also in a primate model mimicking the clinical setting. The effect of LMW-DS was compared with that of heparin, which is routinely used in clinical islet transplantation. LMW-DS was proved to be

Table 1. Summary of the morphological score (as depicted in Fig. 3) of the islets grafts in recipient monkeys M5 and M7–M10

Monkey number	Treatment	Integrity	Thrombus	Cell infiltration	Percentage of score 0 ^a (%)	APTT at 24 h after transplantation (s)
M5	LMW-DS, n = 113	0.66 ± 0.04 ^b	0.26 ± 0.03	0.90 ± 0.07	37.2	101
M7	LMW-DS, n = 134	0.93 ± 0.06	0.52 ± 0.06	1.08 ± 0.06	26.1	66
M8	Heparin, n = 149	1.05 ± 0.05	0.62 ± 0.06	1.17 ± 0.06	20.1	24
P-value ^c		0.13	0.28	0.32	0.23	
M9	LMW-DS, n = 134	0.63 ± 0.05	0.37 ± 0.05	0.85 ± 0.06	44.0	107
M10	Heparin, n = 125	0.95 ± 0.06	0.54 ± 0.06	1.25 ± 0.07	22.4	25
P-value ^d		<0.0001	<0.05	<0.0001	<0.001	

^aPercentage islets with no signs of IBMIR (score 0); ^bValues are expressed as mean ± SEM; ^cP-values for M7 and M8; ^dP-values for M9 and M10. APTT, activated partial thromboplastin time; IBMIR, instant blood-mediated inflammatory reaction; LMW-DS, Low molecular weight dextran sulfate.

Fig. 4. Immunohistochemical staining of porcine islet grafts retrieved 24 h after intraportal xenotransplantation from diabetic monkeys treated with LMW-DS or heparin. The figure shows representative expression of insulin and of CD41 (platelets), CD68 (macrophages), and CD3 (T cells) in the grafts. A summary of all transplanted monkeys is presented in Table 2. Magnification 200×.




Table 2. Summary of the immunohistochemical staining (as depicted in Fig. 4) of the islet grafts in recipient monkeys receiving LMW-DS or heparin

Treatment	CD41	C3c	C9	Neutrophil elastase	CD68	MAC 387	CD56	CD3	CD20	IgG	IgM
LMW-DS (n = 21)	(-)(++) 0.59 ± 0.19 ^a	(-)(++) 0.80 ± 0.37	(-)(+++) 1.50 ± 0.31	(-)(++) 0.42 ± 0.23	(-)(++) 1.31 ± 0.21	(-)(++) 0.90 ± 0.28	(-)(+) 0.10 ± 0.10	(-)(++) 0.63 ± 0.20	(-)(+) 0.20 ± 0.13	(-) 0	(-)(+) 0.25 ± 0.25
Heparin (n = 18)	(-)(+++) 1.60 ± 0.51	(-)(++) 0.63 ± 0.26	(-)(+++) 1.67 ± 0.33	(-)(++) 1.08 ± 0.23	(+)(+++) 2.17 ± 0.11	(+)(+++) 2.11 ± 0.26	(-)(+) 0.22 ± 0.15	(-)(+++) 1.90 ± 0.35	(-)(++) 0.40 ± 0.27	(-) 0	(-)(+++) 0.60 ± 0.24
P value	0.056	0.69	0.71	0.04	0.002	0.01	0.48	0.006	0.83	-	0.36

^aValues are expressed as mean ± SEM. LMW-DS, Low molecular weight dextran sulfate.




Fig. 5. Porcine islets incubated in hirudin-treated plasma for 30 min. The islets were stained for IgG (n = 5), IgM (n = 5), C3b/iC3b (n = 5), C1q (n = 3), C4 (n = 5), C9 (n = 3), and MBL (n = 3). As negative control, an antibody recognizing mouse IgG was used (n = 5). The islets were analyzed by (A) large particle flow cytometry and (B) confocal microscopy. In (C), the deposition of C3b/iC3b on the islet in the absence and presence of Compstatin is presented after analysis by large particle flow cytometry (n = 5; statistical evaluation was performed at 5 and 30 min where n = 7; *P < 0.05 and **P < 0.01).

far more efficient in inhibiting the IBMIR than heparin. These data confirm those of Rood et al. [19] who recently demonstrated longer porcine islet survival in non-human primates treated with LMW-DS.

In this study, both the morphological findings and the measurements in the plasma were similar to those in our previous studies, in which APIs were surrounded by clots and infiltrated by numerous leukocytes immediately after contact with fresh blood seen in the tubing loop model and our small animal model [9]. Most parameters reflecting the IBMIR, i.e. both coagulation and complement cascades, platelet deposition, and infiltration of macrophages and neutrophils, were attenuated in the monkeys treated with LMW-DS compared to

the controls. There was also a tendency that increases in granulocyte count and liver enzymes were attenuated. One control monkey (M6) died of hypoglycemia, suggesting a strong IBMIR. Notably, T-cell infiltration observed in some of the transplanted islet grafts was also effectively suppressed, demonstrating that also the adaptive immune responses are attenuated by LMW-DS.

The effects of LMW-DS on the adaptive immune system may be explained by the effects on complement activation as complement is also of great importance in bridging innate immunity and specific immune responses. In allogeneic whole organ transplantation, C3 is one of the essential factors that trigger rejection in mice [20–22] and humans [23]. It is therefore reasonable to expect that




Fig. 6. Binding of C3b/iC3b to the surface of microtiter wells after incubation with 10% serum in the presence of increasing doses of Compstatin for 30 min at 37 °C. 0 (cross), 10 (triangle), 100 (squares), and 1000 (diamond) mg/l of LMW-DS was present in the wells.

complement activation will trigger a profound adaptive immune response raised against the graft, necessitating an unwarrantedly heavy immunosuppressive regimen. Previous studies support such a hypothesis [4,5].

As shown in Table 1, the islet grafts in M9 that reached an APTT of 107 s 24 h after transplantation, demonstrated well-preserved morphology suggesting that this dose of LMW-DS would be preferable. In a recently performed phase I study in normal individuals, we have shown that this concentration can be reached without an increased risk of bleeding or side effects (manuscript under preparation). This makes treatment with LMW-DS during xenogeneic islet transplantation an attractive alternative. It should be noted that a specific concentration of LMW-DS gives different APTT in blood from different individuals both in vitro and in vivo, probably due to that different allotypes of certain coagulation factors interact with LMW-DS differently.

In our previous studies, we showed that complement activation induced in xenogeneic IBMIR occurs secondarily to coagulation activation; a reaction which is also seen in allogeneic IBMIR [9,24] and which is elicited by chondroitin sulfate released by activated platelets [25]. This explains why complement activation in this study was suppressed in parallel with the reduction of coagulation activation at substantially lower concentrations (15–35 mg/l) of LMW-DS than used in other studies aiming for an inhibition of hyper acute

rejection [26,27]. However, unlike complement activation in the fluid phase, immunohistochemical analyses showed that complement deposition was still seen on the islet grafts in the monkeys treated with LMW-DS. These reactions were analyzed in detail in vitro using large particle flow cytometry and confocal microscopy to be able to clarify the mechanism of activation. The experiments were performed using human plasma to directly translate the findings to clinical islet xenotransplantation. Pig islets incubated in human plasma revealed an almost instantaneous binding of IgM and IgG antibodies and complement components already after 5 min. This rapid activation was completely inhibited by Compstatin. It is possible that the instantaneous insulin dumping in a non-human primate model previously reported by Bennet et al. [3] and also observed in monkey M6 is explained by this antibody-mediated reaction. The severity of this reaction, which was totally abrogated by the recombinant complement inhibitor CR1, is reflected in the fact that the release of insulin corresponded to about 40% of the insulin in the transplanted islets.

One way to fully inhibit complement activation is to increase the dose of LMW-DS, but as shown in our in vitro experiments, doses between 10 and 100 mg/l have only minor effects on complement activation alone. Moreover, higher doses of LMW-DS are likely to give side effects. It is therefore obvious that LMW-DS must be combined with a specific complement inhibitor such as Compstatin to block the immediate destructive immunoglobulin-triggered complement activation found both in vitro and in vivo [10,28]. The in vitro studies show that LMW-DS and Compstatin do not interact in human serum.

Taken together, it is possible to propose a model of how the different components of IBMIR interact in xenogeneic combinations: (1) Immediately when porcine islets come in contact with human blood there is an instantaneous binding of IgG and IgM antibodies to the islet surface which triggers a deleterious complement activation; (2) This is followed by a clotting reaction with accompanying complement activation. Based upon the experimental data presented, LMW-DS combined with a specific complement inhibitor is an attractive alternative to control the detrimental innate immune responses that are postulated to occur in forthcoming intraportal pre-clinical and clinical islet xenotransplantation trials. We are at the moment in progress to produce sufficient amounts of Compstatin to perform studies in the NHP model with Compstatin combined with LMW-DS.

Acknowledgments

This work was supported by National Institute of Health Grants 5U01-A1065192, GM-62134, GM069736, and AI068730, Juvenile Diabetes Research Foundation, 1-2004-18, Swedish Research Council Grants 5647, and 15244.

References

- SHAPIRO AM, LAKEY JR, RYAN EA et al. Islet transplantation in seven patients with type 1 diabetes mellitus using a glucocorticoid-free immunosuppressive regimen. *N Engl J Med* 2000; 343: 230.
- ROOD PP, BUHLER LH, BOTTINO R, TRUCCO M, COOPER DK. Pig-to-nonhuman primate islet xenotransplantation: a review of current problems. *Cell Transplant* 2006; 15: 89.
- BENNET W, SUNDBERG B, LUNDGREN T et al. Damage to porcine islets of Langerhans after exposure to human blood *in vitro*, or after intraportal transplantation to cynomolgus monkeys: protective effects of sCRI and heparin. *Transplantation* 2000; 69: 711.
- BUHLER L, DENG S, O'NEIL J et al. Adult porcine islet transplantation in baboons treated with conventional immunosuppression or a non-myeloablative regimen and CD154 blockade. *Xenotransplantation* 2002; 9: 3.
- CANTAROVICH D, BLANCHO G, POTIRON N et al. Rapid failure of pig islet transplantation in non human primates. *Xenotransplantation* 2002; 9: 25.
- KIRCHHOF N, SHIBATA S, WIJKSTROM M et al. Reversal of diabetes in non-immunosuppressed rhesus macaques by intraportal porcine islet xenografts precedes acute cellular rejection. *Xenotransplantation* 2004; 11: 396.
- CARDONA K, KORBUTT GS, MILAS Z et al. Long-term survival of neonatal porcine islets in nonhuman primates by targeting costimulation pathways. *Nat Med* 2006; 12: 304.
- HERING BJ, WIJKSTROM M, GRAHAM ML et al. Prolonged diabetes reversal after intraportal xenotransplantation of wild-type porcine islets in immunosuppressed nonhuman primates. *Nat Med* 2006; 12: 301.
- GOTO M, JOHANSSON H, MAEDA A, ELGUE G, KORSGREN O, NILSSON B. Low molecular weight dextran sulfate prevents the instant blood-mediated inflammatory reaction induced by adult porcine islets. *Transplantation* 2004a; 77: 741.
- NILSSON B, LARSSON R, HONG J et al. Compstatin inhibits complement and cellular activation in whole blood in two models of extracorporeal circulation. *Blood* 1998; 92: 1661.
- BRANDHORST H, BRANDHORST D, HERING BJ, BRETZEL RG. Significant progress in porcine islet mass isolation utilizing liberase HI for enzymatic low-temperature pancreas digestion. *Transplantation* 1999; 68: 355.
- KISSANE JM, ROBINS E. The fluorometric measurement of deoxyribonucleic acid in animal tissues with special reference to the central nervous system. *J Biol Chem* 1958; 233: 184.
- JOHANSSON H, GOTO M, DUFRANE D et al. Low molecular weight dextran sulfate: a strong candidate drug to block IBMIR in clinical islet transplantation. *Am J Transplant* 2006; 6: 305.
- NILSSON EKDAHL K, NILSSON B, PEKNA M, NILSSON UR. Generation of iC3 at the interface between blood and gas. *Scand J Immunol* 1992; 35: 85.
- MOLLNES TE, RIESENFELD J, GARRED P et al. A new model for evaluation of biocompatibility: combined determination of neoepitopes in blood and on artificial surfaces demonstrates reduced complement activation by immobilization of heparin. *Artif Organs* 1995; 19: 909.
- KATRAGADDA M, MAGOTTI P, SFYROERA G, LAMBRIS JD. Hydrophobic effect and hydrogen bonds account for the improved activity of a complement inhibitor, compstatin. *J Med Chem* 2006; 49: 4616.
- FERNANDEZ LA, HATCH EW, ARMANN B et al. Validation of large particle flow cytometry for the analysis and sorting of intact pancreatic islets. *Transplantation* 2005; 80: 729.
- GOTO M, GROTH CG, NILSSON B, KORSGREN O. Intraportal pig islet xenotransplantation into athymic mice as an *in vivo* model for the study of the instant blood-mediated inflammatory reaction. *Xenotransplantation* 2004b; 11: 195.
- ROOD PP, BOTTINO R, BALAMURUGAN AN et al. Reduction of early graft loss after intraportal porcine islet transplantation in monkeys. *Transplantation* 2007; 83: 202.
- PRATT JR, BASHEER SA, SACKS SH. Local synthesis of complement component C3 regulates acute renal transplant rejection. *Nat Med* 2002; 8: 582.
- CHOWDHURY P, ZHOU W, SACKS SH. Complement in renal transplantation. *Nephron Clin Pract* 2003; 95: c3.
- PENG Q, LI K, PATEL H, SACKS SH, ZHOU W. Dendritic cell synthesis of C3 is required for full T cell activation and development of a Th1 phenotype. *J Immunol* 2006; 176: 3330.
- BROWN KM, KONDEATIS E, VAUGHAN RW et al. Influence of donor C3 allotype on late renal-transplantation outcome. *N Engl J Med* 2006; 354: 2014.
- OZMEN L, EKDAHL KN, ELGUE G, LARSSON R, KORSGREN O, NILSSON B. Inhibition of thrombin abrogates the instant blood-mediated inflammatory reaction triggered by isolated human islets: possible application of the thrombin inhibitor melagatran in clinical islet transplantation. *Diabetes* 2002; 51: 1779.
- HAMAD O, NILSSON EKDAHL K, NILSSON P et al. Complement activation is triggered by chondroitin sulfate released by thrombin receptor activated platelets. *J Thrombosis Haemostasis* 2008, in press.
- FIORANTE P, BANZ Y, MOHACSI PJ et al. Low molecular weight dextran sulfate prevents complement activation and delays hyperacute rejection in pig-to-human xenotransplantation models. *Xenotransplantation* 2001; 8: 24.
- THOMAS H, MAILLET F, LETOURNEUR D, JOZEFONVICZ J, FISCHER E, KAZATCHKINE MD. Sulfonated dextran inhibits complement activation and complement-dependent cytotoxicity in an *in vitro* model of hyperacute xenograft rejection. *Mol Immunol* 1996; 33: 643.
- SOULIKA AM, KHAN MM, HATTORI T et al. Inhibition of heparin/protamine complex-induced complement activation by Compstatin in baboons. *Clin Immunol* 2000; 96: 212.